

690295

No. of Printed Pages : 7

BCS-054

**BACHELOR OF COMPUTER  
APPLICATIONS (BCA) (Revised)**

**Term-End Examination**

**December, 2019**

**BCS-054 : COMPUTER ORIENTED NUMERICAL  
TECHNIQUES**

*Time : 3 Hours*

*Maximum Marks : 100*

---

*Note : Question No. 1 is compulsory. Attempt any  
three more questions from the questions no.  
2 to 5. Use of any calculator is permitted.*

---

1. (a) Find the absolute error and relative error  
in the numbers 432.8 and 0.12584 if four  
digit mantissa is used and chopping is used  
for approximation. 4

(b) Round the following numbers to two  
decimal places : 2

38.21416, 4.3742, 82.375, 2.4869

- (c) For the following two floating point numbers : 3

$$x_1 = 0.5527 \times 10^4$$

and  $x_2 = 0.6243 \times 10^3$

find  $x_1 - x_2$ . The result should be rounded to four decimal digits.

- (d) Find the product of  $x_1$  and  $x_2$  given in Q. No. 1 (c) above. The result should be chopped to four decimal digits. 3

- (e) Find the Newton's forward difference interpolating polynomial for the following data. Hence obtain the value of  $f(x)$  at  $x = 1.5$  : 6

$x$	$f(x)$
1	34
2	60
3	90
4	124
5	162
6	204

- (f) Write the following system of linear equations in matrix form : 2

$$x + 2y + 3z = 14$$

$$x - y = -1$$

$$y + 3z = 11$$

- (g) Solve the following system of linear equations using Gauss-Seidel iterative method : 5

$$x + 6y = 13$$

$$4x - y = 2$$

Perform two iterations, taking  $x = 0$  and  $y = 0$  as the initial values.

- (h) Find an interval in which the following equation has a positive root : 2

$$2x^3 + x^2 - 20x + 12 = 0$$

- (i) Find  $\Delta f$  for the following functions for some  $h > 0$  : 3

(i)  $f(x) = 3x^2$

(ii)  $f(x) = 2x$

- (j) Find the approximate value of  $I = \int_0^1 \frac{dx}{1+x^2}$  using Trapezoidal rule dividing the interval into five equal parts. 10

2. (a) Using an 8-decimal digit floating point representation (4 digits for mantissa, 2 for exponent and 1 each for sign for exponent and sign for mantissa) represent the following numbers in normalised floating point form (using chopping if required): 6

(i) 92752

(ii) -93.231

(iii) -0.0012345

(b) Solve the following system of linear equations by using Gaussian elimination method: 8

$$x_1 - x_2 - x_3 = -3$$

$$2x_1 + 3x_2 + 5x_3 = 7$$

$$x_1 - 2x_3 + 3x_3 = -11$$

(c) Give one example each of the following: 6

(i) Ill conditional problem

(ii) Ordinary differential equation (ODE) of degree 3 and order 2

(iii) A system of inconsistent linear equations in two variables.

3. (a) Consider the initial-value problem : 6

$$y' = 0.2xy, y(1) = 1$$

Use Euler's method to obtain an approximation to  $y(1.2)$  using  $h = 0.1$ .

- (b) Using Lagrange's interpolation formula, find the form of the function  $y(x)$  from the following table. Also compute  $f(3)$  : 7

$x$	$y$
0	6
2	20
5	56

- (c) Write the expressions, one for each, which is obtained by applying each of the following operators to  $f(x)$  for some  $h > 0$  : 4

(i)  $\nabla$

(ii)  $\delta$

(iii)  $\mu$

(iv)  $E$

- (d) Derive the relation between  $\delta$  and  $E$ . 3

4. (a) Solve the following system of linear equations using partial pivoting : 10

$$x + y - 5z = 0$$

$$5x + 2y - z = 18$$

$$2x - 2y + z = 3$$

- (b) Find a real root for the equation  $x^3 + x - 5 = 0$ .

Using Regula-Falsi method, taking  $x$  coordinates of initial points as  $x = 0$  and  $x = 2$ . Perform only two iterations of the method. 7

- (c) Make the Newton's divided difference table for the following data : 3

$x$	$f(x)$
1	10
2	20
4	40
8	80

5. (a) Explain the concept of overflow and underflow in the context of decimal floating point number with the help of *one* example of each. 6

- (b) Find by Newton-Raphson's method, the real root of the equation  $x^2 - 3x + 1 = 0$  taking  $x = 2$  as the starting value. Show three iterations. 7
- (c) Apply Newton's backward difference formula to the data below to obtain a polynomial of degree 4 in  $x$ : 7

$x$	$y$
1	1
2	-1
3	1
4	-1
5	1