

No. of Printed Pages : 6

MCSE-004

MCA (Revised)

Term-End Examination, 2019

MCSE-004 : NUMERICAL AND STATISTICAL COMPUTING

Time : 3 Hours

Maximum Marks : 100

---

**Note :** Question No. 1 is compulsory. Attempt any three questions from the rest. Use of calculator is allowed.

---

1. (a) If 0.333 is the approximate value of  $(1/3)$ , find absolute, relative and percentage error. [3]
- (b) Determine the number of iterations required, to obtain the smallest positive root of  $x^3 - 2x - 5 = 0$ , correct upto two decimal places, when Bisection method is used. [5]
- (c) Solve the following system of equations using Gauss Elimination method : [5]

$$x + 2y + z = 3$$

$$2x + 3y + 3z = 10$$

$$3x - y + 2z = 13$$



(d) ✓ Write probability distribution formula for Binomial distribution and Poisson distribution. [2]

(e) ✓ Determine the value of  $f(x)$  at  $x=4$ , using Lagrange Interpolation formula on the data given below : [5]

x	1.5	3	6
f(x)	2.5	2	20

(f) Calculate the value of  $\int_0^{0.6} e^x dx$ , correct to five significant figures; by using Simpson's (1/3) rule. (Take  $n=6$ ). [5]

(g) If a bank receives on an average  $\lambda = 6$  bad cheques per day. What is the probability that it will receive 4 bad cheques on any given day? [5]

(h) Following data is given for marks in subject A and B of certain examination : [7]

	Subject A	Subject B
Mean Marks	36	85
Standard Deviation	11	8

Given, the coefficient of correlation between A and B =  $\pm 0.66$ .

(i). Determine the two equations of regression.

(ii) Calculate the expected marks in A, corresponding to 75 marks obtained in B.

(i) For  $x = 0.5555 E1$ ;  $y = 0.4545 E1$  and  $z = 0.4535 E1$ , prove that  $x(y - z) \neq xy - xz$ .  
[3]

2. (a) Solve the Quadratic equation  $4x^2 + 8x - 21 = 0$  using two decimal digit arithmetic with rounding, using any two of the following methods : [10]

(i) Bisection method

(ii) Secant method

(iii) Regula Falsi method

(b) Solve the initial value problem  $u' = -2tu^2$ , with  $u(0) = 1$ ,  $h = 0.2$  on the interval  $[0, 1]$ . Use the fourth order classical Runge-Kutta method. [10]

3. (a) Obtain the positive root of the equation  $x^2 - 1 = 0$  by Newton-Raphson method, correct to two decimal places. [8]



- (b) Solve the following system using LU decomposition method : [7]

$$6x_1 - 2x_2 = 14$$

$$9x_1 - x_2 + x_3 = 21$$

$$3x_1 + 7x_2 + 5x_3 = 9$$

- (c) Determine the lowest degree polynomial, which satisfies the following set of values, using forward difference. Also find the polynomial : [5]

x	0	1	2	3	4	5	6	7
f(x)	0	7	26	63	124	215	342	511

4. (a) Calculate the value of the Integral  $\int_4^{5.2} \log x \, dx$  by using Simpson's (1/3) rule and Simpson's (3/8) rule. [10]
- (b) Compute the value of R and  $R^2$  for the data given below, where  $R = S_{xy} / \sqrt{S_{xx} S_{yy}}$ . [5]

Sample Size (i)	12	21	15	1	24
$x_i$	0.96	1.28	1.65	1.84	2.35
$y_i$	138	160	178	190	210
$\hat{y}_i$	138				
$\hat{e}_i$	0				

Regression equation  $\hat{y}_i = 90 + 50x_i$  is used to fill the table where  $\hat{e} = y_i - \hat{y}_i$ .

- (c) Solve the following system of equations by using Jacobi Method, determine the results for two approximations : [5]

$$3x + 4y + 15z = 54.8$$

$$x + 12y + 3z = 39.66$$

$$10x + y - 2z = 7.74$$

5. (a) Write Short notes on the following : [6]

(i) Binomial Distribution

(ii) Poisson Distribution

(iii) Normal Distribution

- (b) Three bags of same type have the following balls : [6]

Bag 1 : 2 Black and 1 White

Bag 2 : 1 Black and 2 White

Bag 3 : 2 Black and 2 White

One of the bag is selected and one ball is drawn. It turns out to be white. What is the probability of drawing a white ball again :

- (i) When the first one is returned/replaced.
- (ii) When the first one is not returned/replaced.

- (c) Calculate the correlation coefficient for the following data : [8]

x :	65	66	67	67	68	69	70	72
y :	67	68	65	68	72	72	69	71

Obtain the equations of lines of regression. Also estimate the value of x for y = 70.

----- X -----